

The Effect of Zinc on the Viability and Seedling Growth of Deteriorated Soybean and Rice Seeds

M.O. Ajala*, N. O. Adedipe** and D. K. Ojo*

*Department of Plant Breeding and Seed Technology, University of Agriculture, P M B 2240, Abeokuta, Ogun State, Nigeria.

** Department of Crop Protection and Environmental Biology, University of Ibadan, Ibadan, Nigeria.

(Accepted 4 July 2003)

ABSTRACT

An investigation was carried out to study the effect of the application of the micro-element Zinc, on the viability and seedling growth of deteriorated soybean (TGX 536-02D) and rice (IR-5) seed. One batch of the seeds was naturally aged and the other subjected to accelerated ageing. Four levels of ZnSO₄ concentration used were 0, 1, 10 and 100 mg/l. For the naturally aged seed, even though there was no significant effect on soybean and rice seed viability for all the concentration levels tested, there was a beneficial carryover effect resulting in significant enhancement of seedling growth in soybean. On seeds subjected to accelerated ageing, 100 mg Zn/l improved soybean seed viability by about 115% and also had promotive effects on seedling growth, notably seedling weight and leaf area by 42% and 25% respectively. The implications of Zn application in the invigoration of seeds with depressed viability were discussed.

Key words: Accelerated ageing, invigoration, rice, seed viability, soybean.

INTRODUCTION

The application of minerals and chemicals to revive seeds of low viability has been the subject of research in the past three decades or so. Both macro- and micro nutrients as well as growth regulators have been employed by many researchers in seed invigoration experiments.

As early as 17th century, it was established that treatment of seeds with salty water protected them from spoilage. The use of chemical fertilizers at the time of sowing was discovered to have an enhancing effect on germination and growth of seedlings (Ovcharov, 1969). It was specifically reported that rye germination was markedly increased by 84% when its seeds were soaked in a 0.05% solution of P₂O₅. The effect was even more dramatic on oats (Ovcharov, 1965).

The performance of plants from aged seeds is known to differ with respect to plant populations (Oladiran, 1994; Oladiran and Agunbiade, 2000).

Iodination of mustard seeds demonstrated an appreciable degree of protection against damage under a variety of storage conditions (Basu and Rudrapal, 1980). In their study, seeds treated with iodine retained a germination of approximately 100% while untreated seeds declined to about 75% germination, presumably as a result of lipid peroxidation.

Microelements have similarly been employed in some seed invigoration studies (Baertuev, 1965;

Kefford *et al.*, 1965; Bekendam and Bruinsma, 1966; Lang, 1970; Adedipe and Ormrod, 1977; Ajala, 1988).

Generally, when microelements are fed to seeds, they do have a very notable influence on many aspects of metabolism (Rogers *et al.*, 1995). Having stored seeds inevitably in an unsuitable environment with its attendant lowered viability, it should be possible to re-invigorate such seeds by chemical application.

The aim of this investigation was therefore to find out the effect of Zinc absorption on seed viability and seedling growth of soybean and rice seeds.

MATERIALS AND METHODS

Soybean (cv. TGX 536-02D) and rice (cv. IR-5) seeds were obtained from the International Institute of Tropical Agriculture (IITA), Ibadan. Ambient storage (26-29°C temperature; 65-80% RH) was carried out for soybean and rice for a period of 2 months and 6 months respectively, and were referred to as 'naturally aged seeds'. Germination percentage was immediately determined before further tests were carried out.

Freshly harvested seeds of the same species and cultivar were procured and also subjected to accelerated ageing conditions (45°C and 95 - 100% RH, 5 days) before chemical application. The concentrations of the chemical were chosen on the bases of specific responses obtained by other workers (Baertuev, 1965; Simakin, 1966; Adedipe, 1973;

Ajala and Adedipe, 1999a; Ajala and Adedipe, 1999 b). Hence, four concentrations used in this study were 0, 1, 10, 1000 mg per litre.

For each batch of soybean and rice seeds, ten seeds were weighed and put in Petri dishes after which 10ml. of Zinc Sulphate was added at 0, 1, 10 and 100 mg/l concentrations. After 8h., they were then transferred into Petri dishes containing distilled water for another 5 days. The number of seeds that germinated was estimated. For seedling growth measurements, after the estimates of the germination percentage, seedlings were transferred into pots containing about 6 kg soil for further growth and observation. On the 10th day, i.e. 15 days after initiation of seedling growth, measurements were taken on length and weight of plumule and radicle as well as leaf weight and total leaf area. Leaf area was estimated by the graphic method.

The experiment was repeated 3 times. The data were subjected to statistical analysis and the Duncan's (1955) New-Multiple Range Tests were used in testing the treatment means where appropriate.

RESULTS

A. Naturally Aged Seeds

Seed Viability

Soybean showed no significant effect for all the Zn concentrations tested, except at 100mg Zn/l where the highest viability was obtained (Fig. 1) corresponding to the highest response in most seedling growth characters (Fig. 2). No significant effect of Zinc was noticed on rice viability (Fig. 1).

Seedling growth responses

There was significant enhancement of seedling growth in soybeans at 10 and 100 mg Zn/l (Fig. 2). With the exception of radicle length and radicle fresh weight in which there were no significant increases, all the other growth parameters recorded significant increases. Similarly, with the exception of radicle fresh weight which was significantly increased at 1mg/l in rice, all other growth parameters were not significantly different from the control (Table. 1).

B. Seeds Subjected to Accelerated Ageing

Seed Viability

Seed viability of soybean was significantly enhanced by 115 % at 100 mg Zn/l (Fig. 3). Rice on the other hand showed a significant reduction in its viability at higher concentrations (10 and 100 mg Zn/l), even though there was an initial significant response at 1 mg/l.

Seedling growth responses

In soybeans, seedling growth parameters, especially plumule fresh weight and leaf area were significantly

enhanced at 100mg Zn/l (Table 2). Its enhancing effect on seedling growth characteristics is worthy of note. Seedling weight was distinctly increased by 42% while leaf area was increased by 25%. Also there were significant differences in radical weight and length occurring at 1 mg/l but not at higher concentrations. In rice, apart from plumule weight which showed a significant increase at 1 mg/l no significant effect was observed for the growth parameters except seedling weight which became depressed at 100 mg Zn/l (Table 2).

Table 1: Effects of Zinc absorption on seedling growth of naturally aged rice seeds.

	Zinc Conc. (mg/l)	Plumule	Radicle
Seedling wt. (g)	0	0.120±0.003	0.13 ± 0.01
	1	0.180±0.010	0.23 ± 0.01
	10	0.110 ± .003	0.10 ± 0.01
	100	0.120±0.002	0.11 ± 0.01
Leaf area (cm ²)	0	0.060 ± .005	9.4 ± 0.4
	1	0.110±0.010	13.5 ± 0.7
	10	0.050±0.005	9.0 ± .12
	100	0.060±0.004	9.0 ± 0.5
Seedling length (cm)	0	17.5 ± 0.5	6.5 ± 0.8
	1	18.0±0.6	7.8 ± 1.5
	10	16.2 ± 0.6	10.5 ± 1.0
	100	15.5 ± 0.5	8.2 ± 0.9

Each figure is a mean of three replicates ± standard error.

DISCUSSION

This study describes the effects of zinc on rice and soybean seed viability and seedling growth following natural and accelerated seed ageing. On naturally aged seed, the microelement (Zn), studied has not been particularly promotive on soybean and rice seed viability, although seedling growth was significantly enhanced in soybeans at 10 and 100 mg Zn/l. It is interesting to note that some chemicals may not influence seed viability but can still exert their influence on subsequent seedling growth. In the studies carried out by Masev and Kutacek (1966), soaking of barley seeds in a solution of ZnSO₄ led to an increased content of growth substances, particularly IAA. However, Bewley and Black (1984) later stated that there is no substantial evidence that IAA participates in germination process and that later events of seedling growth may be controlled by the hormone. Hence, Zn might have enhanced seedling growth, though it may not affect the seed viability of naturally aged soybean. Earlier work carried out by

Table 2: Effects of Zinc absorption on seedling growth in deteriorated* soybean and rice seeds

	Zinc Conc. (mg/l)	Plumule	Radicle
Soybean			
Seedling wt. (g)	0	0.63 ± .03	0.28 ± .02
	1	0.62 ± .02	0.40 ± .02
	10	0.65 ± .03	0.38 ± .03
	100	0.79 ± .06	0.40 ± .03
Leaf area (cm ²)	0	0.16 ± .01	12.33 ± .20
	1	0.18 ± .01	13.67 ± 0.58
	10	0.20 ± .03	14.50 ± 0.29
	100	0.23 ± .03	15.47 ± 1.5
Seedling length (cm)	0	12.00 ± 0.58	8.33 ± 0.33
	1	12.00 ± .58	12.00 ± .58
	10	11.33 ± 0.88	11.67 ± 0.33
	100	13.67 ± 0.88	11.0 ± 1.15
Rice			
Seedling wt. (g)	0	0.089 ± 0.012	0.108 ± .20
	1	0.127 ± 0.006	0.151 ± 0.035
	10	0.100 ± 0.005	0.135 ± 0.007
	100	0.048 ± 0.006	0.076 ± 0.003
Leaf area (cm ²)	0	0.044 ± 0.016	6.33 ± 1.39
	1	0.059 ± 0.006	9.40 ± 0.40
	10	0.052 ± 0.005	6.77 ± 1.23
Seedling length (cm)	0	0.032 ± 0.003	5.57 ± 0.38
	1	13.67 ± 0.88	8.33 ± 0.070
	10	15.83 ± 0.60	6.83 ± 0.73
	100	14.33 ± 1.45	7.33 ± 1.33
	100	12.50 ± 1.26	6.17 ± 0.60

*Seeds were subjected to accelerated ageing conditions (95% RH, 45°C for 5 days) before chemicals were applied.

Each value is the mean of 3 replicates ± standard error.

Ajala and Adedipe (1999a) revealed that ascorbate had anti-oxidant properties which were responsible for lowering the rate of lipid auto-oxidation in stored seeds. Hence deteriorated seeds of rice benefited from ascorbic acid treatment at 10 and 100 mg/l and resulted in significant improvement in viability and seedling growth characters.

Seeds subjected to accelerated ageing, Zn at 100 mg/l resulted in increase in seed germination and growth vigor of seedlings (Fig. 3, Table 2). It is pertinent to note that soybean seed which had initially deteriorated in storage had its viability dramatically improved by 115 % (Fig. 2). Besides, seedling growth characters, notably seedling weight and leaf area increased by 42% and 25% respectively (Table 2).

Therefore, the possibility cannot be excluded that Zn may not be particularly effective on fresh soybean viability (Fig. 1), but its functions of the rehabilitation of the substandard soybean seed are hereby established. This is particularly important with respect to seeds that have suffered one form of disaster or another which had resulted in a sudden drop in viability.

Rice viability became depressed at Zn concentrations of 10 and 100 mg/l. Even though most of the seedling parameters were not affected, seedling weight was distinctly lowered at 100mg Zn/l to about 38%. This shows that rice may not have suffered low levels of endogenous auxin and may therefore not be benefited by Zn treatment. On the other hand, the chemical level required for effect may fall outside the range used in this study.

The present study, however, indicates that Zn is implicated in soybean viability. Even though there was no substantial evidence to prove that viability improvement through zinc application was IAA mediated, Zn had been particularly effective in the enhancement of viability and modification of seedling growth behavior of soybean. This is to say that there is no ambiguity to the role of this micronutrient on soybean seed viability. Although no stimulating effect on naturally aged soybean seed viability was reported, its seedling growth was enhanced by Zn nutrition. Further studies should test changes in IAA concentrations in order to determine whether the observed effects of Zn are IAA mediated or not. This is without prejudice to the fact that the application of this microelement (Zn) has proved beneficial in the modification of viability and seedling growth behaviour of soybean seed subjected to accelerated ageing.

REFERENCES

- Adedipe, N.O. 1973. Effects of 2-chloroethyl phosphonic acid on growth, plastid pigments and sucrose translocation in radish. *J. Exp. Bot.* 24: 124-129.
- Ajala, M. O. 1988. *Physiological quality of stored seeds: Environmental effects and chemical regulation*. Unpublished Ph.D. Thesis, University of Ibadan, Ibadan.
- Ajala, M.O and N.O Adedipe. 1999a. The invigorating role of Ascorbic acid in aged rice seeds. *Bioscience Res. Communication* 11 (4): 20-25.
- Ajala M O and N O Adedipe. 1999b. Temporal Variation in seedling vigour and Total Mineral Contents of Corn and Rice seeds during storage. *Nigerian Journal of science* 33(1):13-20.
- Baertuey, A.A. 1965. Effectiveness of presowing treatment of buck wheat seed with




Figure 1. Effect of Zn absorption on the viability of naturally aged soybean and rice seeds




Figure 2. Effect of Zn absorption on the viability of deteriorated soybean and rice seeds

- microelements (Engl. Transl.) *Trudy Burvatsk* 17:45-47.
- Basu, R.N. and A.B. Rudrapal. 1980. Iodination of mustard seeds for the maintenance of vigour and viability. *Indian J. Exptal. Biol.* 18: 492-496.
- Bewley, J.D. and M. Black. 1984. *Physiology and Biochemistry of seeds*. Vol. 3. Springer Verlag, New York, 260p.
- Duncan, D.B., 1955. Multiple range and multiple F-tests. *Biometrics* 11:1-42. Kefford, N.P.; Zwar, J.A. and Bruce, M.I. 1965. Enhancement of lettuce seed germination by some urea derivatives. *Planta* 67 (1): 103-106
- Lang, A. 1970. Gibberellins: Structure and metabolism. *Ann. Rev. Plant Physiol.* 21:537-539 Masev, N. and M. Kutacek. 1966. The effect of Zinc on the biosynthesis of tryptophan, indol auxins and gibberellins in barley. *Biol. Plant Acad. Sci Biochem* 8(2): 142-151.
- Ovcharov, K.E., 1969. *Physiological basis of seed germination* (Transl. From Russian) Amerind Pub. Co New Delhi
- Rogers, M.E., C.L. Noble, G.M. Halloran and M.I. Nicolai, 1995. The effect of aCl on the germination and early seedling growth of white clover (*Trifolium repens*, L.) populations selected for high and low salinity tolerance. *Seed Sci. and Techn.* 23:277-288.
- Simakin, K.K., 1966. Presowing seed treatment as a method for increasing the quality and production of standard apple seedlings. Avtoreferat Kand. Diss., 125p




Figure 3. Effect of Zn absorption on seedling growth of naturally aged soybean and rice seeds